

Urinary Tract Pathogens (in Order of Frequency) - % Susceptible

| Organism | Number of Isolates | Amox clavulanic | Ampicillin | Cefazolin (1) | Ceftazidime | Ceftriaxone | Ciprofloxacin | Fosfomycin | Gentamicin | Meropenem | Nitrofurantoin | Trimethoprim-Sulfamethoxazole |
|---------------------------|--------------------|-----------------|------------|---------------|-------------|-------------|---------------|------------|------------|-----------|----------------|-------------------------------|
| E. coli ^ | 1417 | 81 | 48 | 81 | | 88 | 56 | 97 | 92 | 100 | 96 | 72 |
| Enterococcus species ^^^^ | 398 | | | | | | | | | | | |
| Proteus mirabilis + | 331 | 97 | 78 | 93 | | 99 | 67 | | 99 | 98 | | 76 |
| Klebsiella pneumoniae * | 316 | 96 | | 91 | | 93 | 88 | | 96 | 100 | 63 | 88 |
| Pseudomonas aeruginosa | 142 | | | | 95 | | 80 | | 89 | | | |
| Group B Streptococcus ^^ | 117 | | | | | | | | | | | |

Organism Notes:

* Includes ESBL and AMP-C isolates (6.6% of total Klebsiella pneumoniae isolates identified as ESBL and AMP-C).

^ Includes ESBL and AMP-C isolates (11.9% of total E.coli isolates identified as ESBL and AMP-C).

^^ This isolate is predictably susceptible to Penicillin.

^^^ Clindamycin, Trimethoprim/Sulfamethoxazole and all Cephalosporins are ineffective against Enterococcus species. Enterococcus isolates recovered from urine are generally susceptible to amoxicillin and nitrofurantoin. Susceptibility to Amoxicillin is 98.7% and to Nitrofurantoin is 96.5%

+ Includes ESBL and AMP-C isolates (1.2% of total Proteus mirabilis isolates identified as ESBL and AMP-C).

Antibiotic Notes:

(1) Cefazolin interpretation predicts results for Cephalexin (Keflex) in accordance with CLSI standards for urinary sites only (not systemic).

All Other Specimen Types excluding (Urines and Surveillance) - Organisms in Order of Frequency - % Susceptible

| Organism | Number of Isolates | Cefazolin | Ceftazidime | Ciprofloxacin | Clindamycin | Cloxacillin | Erythromycin | Gentamicin | Tetracycline (2) | Trimethoprim-Sulfamethoxazole |
|---------------------------|--------------------|-----------|-------------|---------------|-------------|-------------|--------------|------------|------------------|-------------------------------|
| Staphylococcus aureus ^^^ | 491 | 63 | | | 54 | 63 | 43 | | 99 | 100 |
| Pseudomonas aeruginosa | 173 | | 95 | 80 | | | | 97 | | |
| Group B Streptococcus ^^ | 40 | | | | | | | | | |

Organism Notes:

^^ This isolate is predictably susceptible to Penicillin.

^^^ Includes Methicillin Resistant S.aureus (MRSA). MRSA is resistant to all B-Lactams (penicillins, cephalosporins, B-lactam/B-lactamase inhibitor combinations, and carbapenems). MRSA constitutes 35.8% of total Staphylococcus aureus isolates identified.






Antibiotic Notes:

(2) Organisms that are susceptible to Tetracycline are also considered susceptible to Doxycycline.

General Notes:

Antibiogram results, patient risk factors for resistant organisms, and resistance epidemiology should be considered together to help guide empiric treatment of initial infections. Treatment should be re-evaluated as additional information from culture and sensitivity become available.

Calculation of results based on first isolate per patient.

| | |
|------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------|
|  | 90-100% of isolates are susceptible to the antibiotic indicated (GOOD CHOICE) |
|  | 21-89% of isolates are susceptible to the antibiotic indicated (INTERMEDIATE CHOICE) |
|  | 0-20% of isolates are susceptible to the antibiotic indicated (POOR CHOICE) |
|  | Value based on < 30 isolates. Statistical comparison on results with less than 30 isolates is unreliable. n = # of isolates tested. |
|  | Antibiotic susceptibility testing is not typically performed on the organism. |