

Urinary Tract Pathogens (in Order of Frequency) - % Susceptible

| Organism | Number of Isolates | Amox clavulanic | Ampicillin | Cefazolin (1) | Ceftriaxone | Ciprofloxacin | Fosfomycin | Gentamicin | Meropenem | Nitrofurantoin | Trimethoprim-Sulfamethoxazole |
|----------------------------------|--------------------|-----------------|-------------|---------------|-------------|---------------|------------|-------------|--------------|----------------|-------------------------------|
| E. coli ^ | 12600 | 89 | 60 | 88 | 91 | 71 | 99 | 93 | 100 | 98 | 79 |
| Enterococcus species ^^^^ | 3935 | | | | | | | | | | |
| Group B Streptococcus ^^ | 2581 | | | | | | | | | | |
| Klebsiella pneumoniae * | 2004 | 97 | | 93 | 95 | 87 | | 99 | 100 | 27 | 91 |
| Proteus mirabilis + | 837 | 98 n:139 | 87 n:835 | 96 n:835 | 99 n:835 | 92 n:835 | | 94 n:835 | 100 n:835 | n:835 | 89 n:835 |
| Staphylococcus saprophyticus ^^^ | 389 | | | | | | | | | | |

Organism Notes:

* Includes ESBL and AMP-C isolates (4.8% of total Klebsiella pneumoniae isolates identified as ESBL and AMP-C).

^ Includes ESBL and AMP-C isolates (8.4% of total E.coli isolates identified as ESBL and AMP-C).

^^ This isolate is predictably susceptible to Penicillin.

^^^ Acute and uncomplicated urinary tract infections due to Staphylococcus saprophyticus will respond to commonly used antibiotics including Nitrofurantoin, Trimethoprim-Sulfamethaxazole and Fluoroquinolones.

^^^^ Clindamycin, Trimethoprim/Sulfamethoxazole and all Cephalosporins are ineffective against Enterococcus species. Enterococcus isolates recovered from urine are generally susceptible to amoxicillin and nitrofurantoin. Susceptibility to Amoxicillin is 98.7% and to Nitrofurantoin is 96.5%

+ Includes ESBL and AMP-C isolates (0.7% of total Proteus mirabilis isolates identified as ESBL and AMP-C).

Antibiotic Notes:

(1) Cefazolin interpretation predicts results for Cephalexin (Keflex) in accordance with CLSI standards for urinary sites only (not systemic).

All Other Specimen Types excluding (Urines and Surveillance) - Organisms in Order of Frequency - % Susceptible

| Organism | Number of Isolates | Cefazolin | Ceftazidime | Ciprofloxacin | Clindamycin | Cloxacillin | Erythromycin | Gentamicin | Tetracycline (2) | Trimethoprim-Sulfamethoxazole |
|---------------------------|--------------------|-----------|-------------|---------------|-------------|-------------|--------------|-------------|------------------|-------------------------------|
| Staphylococcus aureus ^^^ | 1119 | 86 | | | 76 | 86 | 64 | | 94 | 99 n:997 |
| Group A Streptococcus ^^ | 470 | | | | | | | | | |
| Pseudomonas aeruginosa | 317 | | 95 n:316 | 88 n:316 | | | | 98 n:316 | | |
| Group B Streptococcus ^^ | 136 | | | | | | | | | |

Organism Notes:

^^ This isolate is predictably susceptible to Penicillin.

^^^ Includes Methicillin Resistant S.aureus (MRSA). MRSA is resistant to all B-Lactams (penicillins, cephalosporins, B-lactam/B-lactamase inhibitor combinations, and carbapenems). MRSA constitutes 12.4% of total Staphylococcus aureus isolates identified.

Antibiotic Notes:

(2) Organisms that are susceptible to Tetracycline are also considered susceptible to Doxycycline.

General Notes:

Antibiogram results, patient risk factors for resistant organisms, and resistance epidemiology should be considered together to help guide empiric treatment of initial infections. Treatment should be re-evaluated as additional information from culture and sensitivity become available.

Calculation of results based on first isolate per patient.

| | |
|--|---|
| | 90-100% of isolates are susceptible to the antibiotic indicated (GOOD CHOICE) |
| | 21-89% of isolates are susceptible to the antibiotic indicated (INTERMEDIATE CHOICE) |
| | 0-20% of isolates are susceptible to the antibiotic indicated (POOR CHOICE) |
| | Value based on < 30 isolates. Statistical comparison on results with less than 30 isolates is unreliable. n = # of isolates tested. |
| | Antibiotic susceptibility testing is not typically performed on the organism. |