

## Urinary Tract Pathogens (in Order of Frequency) - % Susceptible

Organism	Number of Isolates	Amox clavulanic	Ampicillin	Cefazolin (1)	Ceftazidime	Ceftriaxone	Ciprofloxacin	Fosfomycin	Gentamicin	Meropenem	Nitrofurantoin	Trimethoprim-Sulfamethoxazole
E. coli ^	239	74	59	78		80	62	92	87	100	96	78
Enterococcus species ^^^^	92											
Klebsiella pneumoniae *	91	80		79		79	74		98	100	48	83
Proteus mirabilis +	33	100	97	97		100	94		97	100		94
Pseudomonas aeruginosa	30				93		80					
Group B Streptococcus ^^	<30											

### Organism Notes:

\* Includes ESBL and AMP-C isolates ( 18.7% of total Klebsiella pneumoniae isolates identified as ESBL and AMP-C ).

^ Includes ESBL and AMP-C isolates ( 19.7% of total E.coli isolates identified as ESBL and AMP-C ).

^^ This isolate is predictably susceptible to Penicillin.

^^^ Clindamycin, Trimethoprim/Sulfamethoxazole and all Cephalosporins are ineffective against Enterococcus species. Enterococcus isolates recovered from urine are generally susceptible to amoxicillin and nitrofurantoin. Susceptibility to Amoxicillin is 97.1% and to Nitrofurantoin is 96.9%

+ Includes ESBL and AMP-C isolates ( 0.0% of total Proteus mirabilis isolates identified as ESBL and AMP-C ).

### Antibiotic Notes:

(1) Cefazolin interpretation predicts results for Cephalexin (Keflex) in accordance with CLSI standards for urinary sites only (not systemic).

## All Other Specimen Types excluding (Urines and Surveillance) - Organisms in Order of Frequency - % Susceptible

Organism	Number of Isolates	Cefazolin	Ceftazidime	Ciprofloxacin	Clindamycin	Cloxacillin	Erythromycin	Tetracycline (2)	Trimethoprim-Sulfamethoxazole
Staphylococcus aureus ^^^	126	64 n:125			78 n:125	64 n:125	54 n:125	97 n:125	100 n:125
Pseudomonas aeruginosa	45		96	89					
Group B Streptococcus ^^	<30								

### Organism Notes:

^^ This isolate is predictably susceptible to Penicillin.

^^^ Includes Methicillin Resistant S.aureus (MRSA). MRSA is resistant to all B-Lactams (penicillins, cephalosporins, B-lactam/B-lactamase inhibitor combinations, and carbapenems). MRSA constitutes 36.5% of total Staphylococcus aureus isolates identified.

### Antibiotic Notes:

(2) Organisms that are susceptible to Tetracycline are also considered susceptible to Doxycycline.

### General Notes:

Antibiogram results, patient risk factors for resistant organisms, and resistance epidemiology should be considered together to help guide empiric treatment of initial infections. Treatment should be re-evaluated as additional information from culture and sensitivity become available.

Calculation of results based on first isolate per patient.

<span style="background-color: #00FF00; width: 15px; height: 10px; display: inline-block;"></span>	90-100% of isolates are susceptible to the antibiotic indicated (GOOD CHOICE)
<span style="background-color: #FFFF00; width: 15px; height: 10px; display: inline-block;"></span>	21-89% of isolates are susceptible to the antibiotic indicated (INTERMEDIATE CHOICE)
<span style="background-color: #FF0000; width: 15px; height: 10px; display: inline-block;"></span>	0-20% of isolates are susceptible to the antibiotic indicated (POOR CHOICE)
<span style="background-color: #CCCCCC; width: 15px; height: 10px; display: inline-block;"></span>	Value based on < 30 isolates. Statistical comparison on results with less than 30 isolates is unreliable. n = # of isolates tested.
<span style="background-color: #808080; width: 15px; height: 10px; display: inline-block;"></span>	Antibiotic susceptibility testing is not typically performed on the organism.